

TWISTED HOCHSCHILD HOMOLOGY AND MACLANE HOMOLOGY

TEIMURAZ PIRASHVILI

ABSTRACT. We prove that $H_i(A, \Phi(A)) = 0$, $i > 0$. Here A is a commutative algebra over the prime field $\mathbb{F}_p$ of characteristic $p > 0$ and $\Phi(A)$ is A considered as a bimodule, where the left multiplication is the usual one, while the right multiplication is given via Frobenius endomorphism and $H_\bullet$ denotes the Hochschild homology over $\mathbb{F}_p$. This result has implications in MacLane homology theory. Among other results, we prove that $HML_\bullet(A, T) = 0$, provided A is an algebra over a field K of characteristic $p > 0$ and T is a strict homogeneous polynomial functor of degree d with $1 < d < \text{Card}(K)$.

1. INTRODUCTION

In this short note we study Hochschild and MacLane homology of commutative algebras over the prime field $\mathbb{F}_p$ of characteristic $p > 0$. Let us recall that MacLane homology is isomorphic to the topological Hochschild homology [13] and to the stable K -theory as well [5].

Let A be a commutative algebra over the prime field $\mathbb{F}_p$ of characteristic $p > 0$ and let $\Phi(A)$ be denote A - A -bimodule, which is A as a left A -module, while the right multiplication is given via Frobenius endomorphism. We prove that the Hochschild homology vanishes $H_i(A, \Phi(A)) = 0$, $i > 0$. The proof makes use a simple result on homotopy groups of simplicial rings, which says that if $R_\bullet$ is a simplicial ring such that all rings involved in $R_\bullet$ satisfy $x^m = x$, $m \geq 2$ identity then $\pi_i(R_\bullet) = 0$ for all $i > 0$. These results has implications in MacLane homology theory. We extend the computation of Franjou-Lannes and Schwartz [4] of MacLane (co)homology of finite fields with coefficients in symmetric S^d and divided powers Γ^d to arbitrary commutative $\mathbb{F}_p$ -algebras, provided that $d > 1$. As a consequence of our computations we show that $HML_\bullet(A, T) = 0$, provided T is a strict homogeneous polynomial functor of degree $d > 1$ and A is an algebra over a field K of characteristic $p > 0$ with $\text{Card}(K) > d$.

Many thanks to referee for his valuable comments and to Petter Andreas Bergh for his remarks.

2. WHEN IT IS TOO EASY TO COMPUTE HOMOTOPY GROUPS

It is well-known that the homotopy groups of a simplicial abelian group $(A_\bullet, \partial_\bullet, s_\bullet)$ can be computed as the homology of the normalized chain complex $(N_\bullet(A_\bullet), d)$, where

$$N_n(A_\bullet) = \{x \in A_n \mid \partial_i(x) = 0, i > 0\}$$

* Partially supported by the grant from MEC, MTM2006-15338-C02-02 (European FEDER support included).

and the boundary map $N_n(A_\bullet) \rightarrow N_{n-1}(A_\bullet)$ is induced by ∂_0 . Our first result shows that if $A_\bullet$ has a simplicial ring structure and the rings involved in $A_\bullet$ satisfy extra conditions then homotopy groups are zero in positive dimensions. This fact is an easy consequence of the following result which is probably well-known

Lemma 1. Let $R_\bullet$ be a simplicial object in the category of not necessarily associative rings and let $x, y \in N_n(R_\bullet)$ be two elements. Assume $n > 0$ and x is a cycle. Then the cycle $xy \in N_n(R_\bullet)$ is a coboundary.

Proof. Consider the element

$$z = s_0(xy) - s_1(x)s_0(y).$$

Then we have

$$\partial_0(z) = xy - (s_0\partial_0(x))y = xy.$$

Moreover,

$$\partial_1(z) = xy - xy = 0,$$

We also have

$$\partial_2(z) = (s_0\partial_1(x))(s_0\partial_1(y)) - x(s_0\partial_1(y)) = 0.$$

Similarly for all $i > 2$ we have

$$\partial_i(z) = (s_0\partial_{i-1}(x))(s_0\partial_{i-1}(y)) - (s_1\partial_{i-1}(x))(s_0\partial_{i-1}(y)) = 0.$$

Hence z is an element of $N_{n+1}(R_\bullet)$ with $\partial(z) = xy$. $\square$

Corollary 2. Let $R_\bullet$ be a simplicial ring. If the rings involved in $R_\bullet$ satisfy $x^m = x$ identity for $m \geq 2$, then

$$\pi_n(R_\bullet) = 0, \quad n > 0.$$

Proof. Take a cycle $x \in N_n(R_\bullet)$, $n > 0$. Then the class of $x = xx^{m-1}$ in $\pi_n(R_\bullet)$ is zero. $\square$

Remark. A more general fact is true. Let $\mathbf{T}$ be a pointed algebraic theory [15] and let $X_\bullet$ be a simplicial object in the category of $\mathbf{T}$ -models [15]. Then $\pi_1(X_\bullet)$ is a group object in the category of $\mathbf{T}$ -models, while $\pi_i(X_\bullet)$ are abelian group objects in the category of $\mathbf{T}$ -models for all $i > 1$. Thus $\pi_i(X_\bullet) = 0$, $i \geq 1$ provided all group objects are trivial. This is what happens for the category of rings satisfying the identity $x^m = x$, $m \geq 2$. Another interesting case is the category of Heyting algebras [3].

3. HOCHSCHILD HOMOLOGY WITH TWISTED COEFFICIENTS

In what follows the ground field is the prime field $\mathbb{F}_p$ of characteristic $p > 0$. All algebras are taken over $\mathbb{F}_p$ and they are assumed to be associative. For an algebra R and an R - R -bimodule B we let $H_\bullet(R, B)$ and $H^\bullet(R, B)$ be the Hochschild homology and cohomology of R with coefficients in B . Let us recall that

$$H_\bullet(R, B) = \text{Tor}_\bullet^{R \otimes R^{op}}(R, B)$$

and

$$H^\bullet(R, B) = \text{Ext}_{R \otimes R^{op}}^\bullet(R, B).$$

Moreover, we let $C_\bullet(R, B)$ be the standard simplicial vector space computing Hochschild homology

$$\pi_\bullet(C_\bullet(R, B)) \cong H_\bullet(R, B).$$

Recall that $C_n(R, B) = B \otimes R^{\otimes n}$, while

$$\partial_0(b, r_1, \dots, r_n) = (br_1, \dots, r_n),$$

$$\partial_i(b, r_1, \dots, r_n) = (b, r_1, \dots, r_i r_{i+1}, \dots, r_n), \quad 0 < i < n$$

and

$$\partial_n(b, r_1, \dots, r_n) = (r_n b, r_1, \dots, r_{n-1}).$$

Here $b \in B$ and $r_1, \dots, r_n \in R$.

Let $n > 1$ be a natural number and let A be a commutative $\mathbb{F}_p$ -algebra. The Frobenius homomorphism gives rise to the functors Φ^n from the category of A -modules to the category of A - A -bimodules, which are defined as follows. For an A -module M the bimodule $\Phi^n(M)$ coincides with M as a left A -module, while the right A -module structure on $\Phi^n(M)$ is given by

$$ma = a^{p^n} m, \quad a \in A, m \in M.$$

Having A - A -bimodule $\Phi^n(M)$ we can consider the Hochschild homology $H_\bullet(A, \Phi^n(M))$. In this section we study these homologies. In order to state our results we need some notation. We let $\psi^n(A)$ be the quotient ring $A/(a - a^{p^n})$, $n \geq 1$ which is considered as an A -module via the quotient map $A \twoheadrightarrow \psi^n(A)$. Thus ψ^n is the left adjoint of the inclusion of the category of commutative $\mathbb{F}_p$ -algebras with identity $x^m = x, m = p^n$ to the category of all commutative $\mathbb{F}_p$ -algebras.

Example 3. Let $n > 1$. If K is a finite field with $q = p^d$ element then $\psi^n(K) = K$ if $n = dt$, $t \in \mathbb{N}$ and $\psi^n(K) = 0$ if $n \neq dt$, $t \in \mathbb{N}$.

Lemma 4. Let A is a commutative algebra over a field K of characteristic $p > 0$ with $\text{Card}(K) > p^n$. Then $\psi^n(A) = 0$, $n > 1$.

Proof. By assumption there exists $k \in K$ such that $k^{p^n} - k$ is an invertible element of K . It follows then that the elements of the form $a^{p^n} - a$ generates whole A . $\square$

Theorem 5. Let A be a commutative $\mathbb{F}_p$ -algebra and $n > 1$. Then

$$H_i(A, \Phi^n(A)) = 0$$

for all $i > 0$ and

$$H_0(A, \Phi^n(A)) \cong \psi^n(A).$$

Proof. The proof consists of three steps.

Step 1. The theorem holds if $A = \mathbb{F}_p[x]$. In this case we have the following projective resolution of A over $A \otimes A = \mathbb{F}_p[x, y]$:

$$0 \rightarrow \mathbb{F}_p[x, y] \xrightarrow{\eta} \mathbb{F}_p[x, y] \xrightarrow{\epsilon} \mathbb{F}_p[x] \rightarrow 0.$$

Here $\epsilon(x) = \epsilon(y) = x$ and η is induced by multiplication by $(x - y)$. Hence for any A - A -bimodule B , we have $H_i(A, B) = 0$ for $i > 1$ and

$$H_0(A, B) \cong \text{Coker}(u) \text{ and } H_1(A, B) \cong \text{Ker}(u),$$

where $u : B \rightarrow B$ is given by $u(b) = xb - bx$. If $B = \Phi^n(\mathbb{F}_p[x])$, then $u : \mathbb{F}_p[x] \rightarrow \mathbb{F}_p[x]$ is the multiplication by $(x^{p^n} - x)$ and we obtain $H_1(A, \Phi^n(A)) = 0$ and $H_0(A, \Phi^n(A)) = \psi^n(A)$.

Step 2. The theorem holds if A is a polynomial algebra. Since Hochschild homology commutes with filtered colimits it suffices to consider the case when $A = \mathbb{F}_p[x_1, \dots, x_d]$. By the Künneth theorem for Hochschild homology (see [10, Theorem X.7.4]) we have $H_\bullet(A, \Phi^n(A)) = H_\bullet(\mathbb{F}[x], \Phi^n(\mathbb{F}[x]))^{\otimes d}$ and the result follows.

Step 3. The theorem holds for arbitrary A . We use the same method as used in the proof of [9, Theorem 3.5.8]. First we choose a simplicial commutative algebra $L_\bullet$ such that each L_n is a polynomial algebra, $n \geq 0$ and $\pi_i(L_\bullet) = 0$ for all $i > 0$, $\pi_0(L_\bullet) = A$. Such a resolution exist thanks to [14]. Now consider the bisimplicial vector space $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$. The s -th horizontal simplicial vector space is the simplicial vector space $L_\bullet^{\otimes s+1}$. By the Eilenberg-Zilber-Cartier and Künneth theorems it has zero homotopy groups in positive dimensions and $\pi_0(L_\bullet^{\otimes s+1}) = A^{\otimes s+1}$. On the other hand the t -th vertical simplicial vector space of $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$ is isomorphic to the Hochschild complex $C_\bullet(L_t, \Phi^n(L_t))$ which has zero homology in positive dimensions by the previous step. Hence both spectral sequences corresponding to the bisimplicial vector space $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$ degenerate and we obtain the isomorphism

$$H_\bullet(A, \Phi^n(A)) \cong \pi_\bullet(\psi^n(L_\bullet)).$$

Now we can use Lemma 2 to finish the proof. $\square$

Corollary 6. Let A be a commutative $\mathbb{F}_p$ -algebra, M be an A -module and $n > 1$. Then there exist functorial isomorphisms

$$H_\bullet(A, \Phi^n(M)) \cong \text{Tor}_\bullet^A(\psi^n(A), M), \quad n \geq 0$$

and

$$H^\bullet(A, \Phi^n(M)) \cong \text{Ext}_A^\bullet(\psi^n(A), M), \quad n \geq 0.$$

In particular, if A is a commutative algebra over a field K of characteristic $p > 0$ with $\text{Card}(K) > p^n$, then

$$H_\bullet(A, \Phi^n(M)) = 0 = H^\bullet(A, \Phi^n(M)).$$

Proof. Observe that $C_\bullet(A, \Phi^n(A))$ is a complex of left A -modules. By Theorem 5 it is a free-resolution of $\psi^n(A)$ in the category of A -modules. Hence it suffices to note that

$$C_\bullet(A, \Phi^n(M)) \cong M \otimes_A C_\bullet(A, \Phi^n(A)),$$

$$C^\bullet(A, \Phi^n(M)) \cong \text{Hom}_A(C_\bullet(A, \Phi^n(A)), M),$$

where C^* denotes the standard complex for Hochschild cohomology. The last assertion follows from Lemma 4. $\square$

Example 7. It follows for instance that $H^i(A, \Phi^n(M)) = 0$, $i > 0$, provided M is an injective A -module and $n > 1$. In particular $H^i(A, \Phi^n(A)) = 0$ if A is a self-injective algebra. On the other hand if $A = \mathbb{F}_p[x_1, \dots, x_d]$ then $H^i(A, \Phi^n(A)) = 0$, $i \neq d$, $n > 1$ and $H^d(A, \Phi^n(A)) = \psi^n(A)$, $n > 1$.

4. APPLICATION TO MACLANE COHOMOLOGY

We recall the definition of MacLane (co)homology. For an associative ring R we let $\mathbf{F}(R)$ be the category of finitely generated free left R -modules. Moreover, we let $\mathcal{F}(R)$ be the category of all covariant functors from the category $\mathbf{F}(R)$ to the category of all R -modules. The category $\mathcal{F}(R)$ is an abelian category with enough projective and injective objects. By definition [8] the *MacLane cohomology* of R with coefficient in a functor $T \in \mathcal{F}(R)$ is given by

$$\mathrm{HML}^\bullet(R, T) := \mathrm{Ext}_{\mathcal{F}(R)}^\bullet(I, T),$$

where $I \in \mathcal{F}(R)$ is the inclusion of the category $\mathbf{F}(R)$ into the category of all left R -modules. One defines MacLane homology in a dual manner (see [13, Proposition 3.1]). For an R - R -bimodule B , one considers the functor $B \otimes_R (-)$ as an object of the category $\mathcal{F}(R)$. For simplicity we write $\mathrm{HML}_\bullet(R, B)$ instead of $\mathrm{HML}_\bullet(R, B \otimes_R (-))$. There is a binatural transformation

$$\mathrm{HML}_\bullet(R, B) \rightarrow H_\bullet(R, B)$$

which is an isomorphism in dimensions 0 and 1.

In the rest of this section we consider MacLane (co)homology of commutative $\mathbb{F}_p$ -algebras.

Lemma 8. For any commutative $\mathbb{F}_p$ -algebra A one has an isomorphism

$$\mathrm{HML}_{2i}(A, \Phi^n(A)) = \psi^n(A), \quad i \geq 0, n > 1,$$

and

$$\mathrm{HML}_{2i+1}(A, \Phi^n(A)) = 0, \quad i \geq 0, n > 1.$$

Proof. According to [12, Proposition 4.1] there exists a functorial spectral sequence

$$E_{pq}^2 = H_p(A, \mathrm{HML}_q(\mathbb{F}_p, B)) \implies \mathrm{HML}_{p+q}(A, B).$$

Here B is an A - A -bimodule. By the well-known computation of Breen [2], Bökstedt [1] (see also [4]) we have

$$\mathrm{HML}_{2i}(\mathbb{F}_p, B) = B$$

and

$$\mathrm{HML}_{2i+1}(\mathbb{F}_p, B) = 0.$$

Now we put $B = \psi^n(A)$ and use Theorem 5 to get $E_{pq}^2 = 0$ for all $p > 0$. Hence the spectral sequence degenerates and the result follows. $\square$

We now consider MacLane cohomology with coefficients in strict polynomial functors [6]. Let us recall that the strict homogeneous polynomial functors of degree d form an abelian category $\mathcal{P}_d(A)$ and there exist an exact functor $i : \mathcal{P}_d(A) \rightarrow \mathcal{F}(A)$ [7]. For an object $T \in \mathcal{P}_d(A)$ we write $\mathbf{HML}_\bullet(A, T)$ instead of $\mathbf{HML}_\bullet(A, i(T))$. Projective generators of the category $\mathcal{P}_d$ are tensor products of the divided powers, while the injective cogenerators are symmetric powers. Let us recall that the d -th divided power functor $\Gamma^d \in \mathcal{F}(A)$ and d -th symmetric functors S^n are defined by

$$\Gamma^d(M) = (M^{\otimes d})^{\Sigma_d}, \quad S^n(M) = (M^{\otimes d})_{\Sigma_d}.$$

Here tensor products are taken over A , Σ_d is the symmetric group on d -letters, which acts on the d -th tensor power by permuting of factors, $M \in \mathbf{F}(A)$ and X^G (resp. X_G) denotes the module of invariants (resp. coinvariants) of a G -module X , where G is a group.

For a functor $T \in \mathcal{F}(A)$ we let $\tilde{T} \in \mathcal{F}(\mathbb{F}_p)$ be the functor defined by

$$\tilde{T}(V) = T(V \otimes A)$$

According to [13, Theorem 4.1] the groups $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$ have an A - A -bimodule structure. The left action comes from the fact that T has values in the category of left A -modules, while the right action comes from the fact that T is defined on $\mathbf{F}(A)$. In particular it uses the action of T on the maps $l_a : X \rightarrow X$, where $a \in A$, $X \in \mathbf{F}(A)$ and l_a is the multiplication on a . Since $T(l_a) = l_{a^d}$ if T is a strict homogeneous polynomial functor of degree d [6], the bimodule $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$ is of the form $\Phi^n(M)$ provided $d = p^n$.

Theorem 9. Let $d > 1$ be an integer and let A be a commutative $\mathbb{F}_p$ -algebra. Then $\mathbf{HML}_\bullet(A, \Gamma^d) = 0$ if d is not a power of p . If $d = p^n$ and $n > 0$, then

$$\mathbf{HML}_i(A, \Gamma^d) = 0 \quad \text{if } i \neq 2p^nt, t \geq 0$$

and

$$\mathbf{HML}_i(A, \Gamma^d) = \psi^n(A) \quad \text{if } i = 2p^nt, t \geq 0.$$

In particular $\mathbf{HML}_\bullet(A, \Gamma^d) = 0$ provided A is an algebra over a field K of characteristic $p > 0$ with $\text{Card}(K) > d$.

Proof. According to [13, Theorem 4.1], [12] there exists a functorial spectral sequence:

$$E_{pq}^2 = \mathbf{H}_p(A, \mathbf{HML}_q(\mathbb{F}_p, \tilde{T})) \implies \mathbf{HML}_{p+q}(A, T).$$

For $T = \Gamma_A^n$ one has $\tilde{T} = \Gamma_{\mathbb{F}_p}^n \otimes A$. Here we used the notation Γ_A^n in order to emphasize the dependence on the ring A . By the result of Franjou, Lannes and Schwartz [4] $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$ vanishes unless $d = p^n$ and $i = 2p^nt$, $t \geq 0$. Moreover in these exceptional cases $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$ equals to $\Phi^n(A)$ (as an A - A -module). Hence the spectral sequence together with Theorem 5 gives the result. $\square$

Corollary 10. Let A be a commutative algebra over a field K of characteristic $p > 0$ with $\text{Card}(K) > d > 1$. If T is a strong homogeneous polynomial functor of degree d . Then

$$\mathbf{HML}_\bullet(A, T) = 0 = \mathbf{HML}^\bullet(A, T).$$

Proof. We already proved that the result is true if T is a divided power. By the well-known vanishing result [11] the result is also true if $T = T_1 \otimes T_2$ with $T_1(0) = 0 = T_2(0)$. Since any object of $\mathcal{P}_d$ has a finite resolution which consists with finite direct sums of tensor products of divided powers [6] the result follows. $\square$

REFERENCES

- [1] M. Bökstedt. The topological Hochschild homology of $\mathbb{Z}$ and $\mathbb{Z}/p$. Unpublished manuscript.
- [2] L. Breen. Extensions du groupe additif. Publ. IHES 48 (1978), 39–125.
- [3] L. Esakia. Heyting algebras. I. Duality theory. (Russian) “Metsniereba”, Tbilisi, 1985. 105 pp.
- [4] V. Franjou, J. Lannes and L. Schwartz. Autour de la cohomologie de MacLane des corps finis. Invent. Math. 115 (1994), no. 3, 513–538.
- [5] V. Franjou, E. Friedlander, T. Pirashvili and L. Schwartz. Rational representations, the Steenrod algebra and functor homology. Panoramas et Synthèses 16. Société Mathématique de France, Paris, 2003. xxii+132 pp.
- [6] E. Friedlander and A. Suslin. Cohomology of finite group schemes over a field. Invent. Math. 127 (1997), no. 2, 209–270.
- [7] V. Franjou, E. Friedlander, A. Scorichenko and A. Suslin. General linear and functor cohomology over finite fields. Ann. of Math. (2) 150 (1999), no. 2, 663–728.
- [8] M. Jibladze and T. Pirashvili. Cohomology of algebraic theories. J. Algebra 137 (1991), no. 2, 253–296.
- [9] J.-L. Loday. Cyclic homology. Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1998.
- [10] S. Mac Lane. Homology. Die Grundlehren der mathematischen Wissenschaften, Bd. 114, Springer. Berlin. 1963 x+422 pp.
- [11] T. Pirashvili. Higher additivizations. Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruz. SSR 91 (1988), 44–54.
- [12] T. Pirashvili. Spectral sequence for MacLane homology. J. of Algebra, 170(1994), 422–427.
- [13] T. Pirashvili and F. Waldhausen. MacLane homology and topological Hochschild homology. J. Pure Appl. Algebra 82 (1992), 81–98.
- [14] D. Quillen. On the (co-) homology of commutative rings. Proc. Sympos. Pure Math., Vol. XVII, New York, 1968. 65–87.
- [15] S. Schwede. Stable homotopy of algebraic theories. Topology 40 (2001), no. 1, 1–41.